

ing from one place to another, packets of letters may be conveyed each way, at the same time, without a possibility of their clashing against each other; and many packets may be conveyed the same way, in the same tube, which can never approach each other, but will all proceed with an uniform motion and equal rapidity to their destination, where, the tube entering an airtight room, the packets will be deposited, and may be delivered or forwarded to the next stage through their proper tubes, commencing in the same room, and their progress can never be impeded by the seasons or the elements."

This proposal did not take practical form until 1854, when Mr. Latimer Clark laid down a 1½ inch lead pipe between the Electric Telegraph Company's Central Station, Lothbury (LY), and the Stock Exchange. An engine exhausted a receiver at LY, and carriers containing the messages were sucked through from the Stock Exchange. The traffic was only required to flow in one direction. In 1858, the system was extended to Mincing lane, and about 1860 Mr. Varley introduced the use of compressed air, so that messages were drawn in one direction by a vacuum, and propelled in the other direction by a plenum. Mr. Clark had previously used a vacuum to work in both directions, a receiver at Mincing lane having been exhausted by the engine at LY, by means of a special pipe laid down in the same trench with the carrier tube.

In 1865 the system was introduced in Paris. Considerable modifications were made in its mode of working. Compressed air was used entirely, and the necessary pressure was obtained by admitting water from the mains into large air reservoirs. This tube served several stations, which were worked intermediately, like a line of railway, or a telegraph current, each station having its own store of power to propel or forward the carrier on to the next place. This mode of obtaining power was found wasteful and expensive, and it has been nearly entirely abandoned in favor of steam working at one end of the circuit.

About the same period (1865) a system was introduced in Berlin by Messrs. Siemens, who used two pipes, laid in the same trench, between the telegraph station and the Bourse, arranged in a circuit, through which a continuous current of air was always kept flowing in the same direction by a double-acting air pump, worked by a steam engine. This last mode of working was tried in London, but it has not proved successful, and it has been abandoned.

It will be seen how closely this system of Siemens' resembles that of Medhurst, and how curiously history works in a circle, for the vision of 1810 has become the stern fact of 1875. In all the places named the pneumatic telegraph has received considerable extension, and it has also been largely introduced in Vienna, where the Parisian system has been adopted.—*Telegraphic Journal.*

ASTRONOMICAL NOTES.

OBSERVATORY OF VASSAR COLLEGE.

The computations and some of the observations in the following notes are from students in the astronomical department. The times of risings and settings of planets are approximate, but sufficiently accurate to enable an ordinary observer to find the objects mentioned. M. M.

Position of the Planets for January, 1875.

Mercury.

Mercury should be looked for after the middle of the month in the southwest, farther north than the point at which the sun is seen to set. It will be in the best position on the 28th, and can be recognized by its white light and by its nearness to Saturn. At this time, Mercury, Saturn, Venus, and Mars can all be seen in the evening.

Venus.

On the 1st of January, Venus rises at 9h. 8m. A. M., and sets at 6h. 32m. P. M. On the 31st, Venus rises at 8h. 45m. A. M., and sets at 7h. 47m. P. M.

Venus will be well seen all through the month, and will be very near Saturn on the 16th, and in conjunction with the moon on the 28th, at which time the planets mentioned above can also be seen.

Mars.

On the 1st of January, Mars will rise at 10h. 55m. A. M., and set at 10h. 21m. P. M. On the 31st, Mars will rise at 10h. 13m. A. M., and set at 9h. 43m. P. M. The moon will be near Mars (apparently) on the 2d of January, and again on the 30th.

Jupiter.

Jupiter is still unfavorably situated for evening observers. On the 1st it rises about 4 in the morning, and sets at 1h. 39m. P. M. On the 31st it rises at 2h. 18m. A. M., and sets at 11h. 56m. A. M.

Saturn.

Saturn rises on the 1st at 9h. 56m. A. M., and sets at 8h. 5m. P. M. On the 31st, Saturn rises at 8h. 7m. A. M., and sets at 6h. 25m. P. M.

Venus and Saturn will have nearly the same apparent position on the 16th, but will be nearer the horizon, and therefore not so conspicuous as were Mars and Saturn in November.

Uranus.

Uranus is in good position, and can be seen with an ordinary telescope. It is among the small stars of *Leo*, rising on the 1st at 7h. 48m. P. M., and on the 31st at 5h. 41m. P. M. On the 31st it comes to the meridian at 12h. 40m. (20m. before one in the morning) and is then 9½° west of Regulus, and 3½° above that star.

Neptune.

Neptune rises at 0h. 35m. P. M. on the 1st, and sets at 1h. 47m. the next morning. On the 31st it rises at 10h. 38m.

A. M., and sets at 11h. 50m. P. M. It cannot be seen without a powerful telescope.

Sun Spots.

The report is from November 17 to December 17 inclusive. From November 5 to November 18 no spots were seen. The photograph of November 18 showed two going off; but before the next picture, November 22, they had disappeared. In the photograph of November 23 there appeared a group of spots on the western limb, a group on the eastern limb, followed by a single one, and, near the center, two very small ones. Clouds prevented photographing on November 23. The pictures of November 24 and November 25 showed only a regular motion of the spots seen on November 22. In the photograph of November 27 there appeared but one large spot on the western limb: the two single ones first observed, near the center, November 22, could no longer be found, and the group which had been seen on the western limb had passed off. The picture of November 29 shows the large spot going off, surrounded by faculae.

Photographing was much interrupted by clouds from November 29 to December 12; but when openings in the clouds allowed observations with the telescope, the sun's disk was seen to be free from spots until December 12, when a small one was seen coming on, but after that date it could not be found. On December 14 a large spot was observed on the very edge. In the photograph of December 17, this spot appeared to be divided into two, and, near the center of the disk, a group of four very small spots was seen, which had not been found before.

Correspondence.

Electricity a Mode of Motion.

To the Editor of the Scientific American:

I take pleasure in briefly meeting the objections of Mr. R. B. West, of Guilford, Conn., to my theory that electricity is nothing more nor less than a motion of the atomic particles of matter. As Mr. West, in his communication, clearly sets forth what may prove a stumbling block to other inquirers, permit me to quote it in full:

"In No. 23, Volume XXXIII, a correspondent advances the theory, if I rightly understand, that electricity is nothing more than motion in the form of an impact or repulsion, communicated from atom to atom, and decreasing in force with the increase in distance from the starting point. This would seem very probable if electricity were capable of being communicated only by direct metallic contact; but on the contrary, it will pass, with comparatively little resistance, through a space made practically devoid of matter; and an inductive disturbance is produced when there could be no possible communication of force. Electricity may possibly be something like an allotropic form of motion, but the definition of an atomic impact alone can scarcely be used."

Instead of entering upon a criticism of Mr. West's statement of facts, namely, that electricity will pass with comparatively little resistance through a (so-called) vacuum, I prefer, for the sake of the argument, to admit that he is correct.

The difficulty seems to consist in a lack of appreciation of matter itself. Mr. West seems to forget that it is impossible to render a space void of matter. What he conceives to be practically *nil*, as perfect a vacuum as can be formed, is really complete materiality, in which the atomic particles of what he conceives to be *nil* are in direct connection with the atomic particles of the metallic conductor, and therefore capable, in greater or less degree, of transmitting a force existing in the metallic conductor. I agree with Mr. West that there could be no transmission of electricity through a space devoid of matter; but it seems to me that, in raising his objection, he should have offered some proof of fact that he has discovered a space void of matter; or, failing to do so, he should have advanced some argument to show that a void is possible, and that, being possible, it is possible to transmit a force through it; otherwise, he is clearly not warranted in denying a material connection between one so-called conductor and another.

The concluding suggestion of Mr. West, that "electricity may possibly be something like an allotropic form of motion," I confess, puts me a long distance at sea. I am quite at a loss, for instance, to conceive of setting something in motion, and then, taking something away, having motion continue on its own account. I would like very much to witness a practical demonstration of that thing; or if that is impossible, to have somebody advance an argument showing that motion may exist independent of matter. It has always seemed to me a requisite of motion that something material shall move.

New York city.

W. E. SAWYER.

The Rattlesnake's Poison.

To the Editor of the Scientific American:

My attention has been attracted by a statement made in your issue of December 4, 1875, page 353. After showing the fallacy of certain stories which have been widely circulated in print, and by word of mouth, which have gained credence, regarding the toxic effect of the spittle of man when administered to venomous reptiles, and relating the incident of the boot, which contained a serpent's fang and was credited with so fatal a record, you state that the inventor of this story did not know that the rattlesnake poison is only active when freshly injected from the poison bag.

The story, is of course, improbable; but the error of your statement is very clearly shown by the following experiments by Mitchell with the venom of that reptile. He says

that "it is difficult to conceive of the singular energy of the venom of the rattlesnake without carefully conducted experimental research, or of the tenacity with which its powers are preserved in the presence of violent chemical reagents and extremes of heat and cold. The dried venom retained its potency after two years of climatic changes; nor was its action in any degree changed by strong sulphuric and hydrochloric acids, ammonia, chlorine water, soda, or potassa. Freezing or prolonged boiling in no way impaired its deadly qualities." He used the venom after five years' keeping, and found it uninjured.

Dr. Weir Mitchell's reports of his exhaustive researches with this virulent body are richly worth perusal, showing as he does the precise manner in which it is so swiftly and fatally transmitted through the serous tissues, and conclusively settling the fact that the serpent cannot inoculate itself, a point which was for a long time disputed. His reports on this subject may be found in the *Smithsonian Contributions*, 1860, and the *New York Medical Journal*, January, 1868.


New York city.

HENRY S. WELCOME.

[For the Scientific American.]

SEEING DISTANT OBJECTS FROM ELEVATIONS.

A correspondent mentions that it is proposed to build, in Fairmount Park, Philadelphia, adjacent to the Centennial buildings, a tower 750 feet high, and asks if it is true that, from the top of this tower, New York city, 90 miles distant, can be seen. According to the rule that the horizon dips 8 feet for every mile, which, for 90 miles, would be a dip of 8 × 90, or 720 feet, a tower of 750 feet high would be taller than necessary. He does, however, doubt if this rule is correct, and this doubt is well founded. If the rule were correct, we could see from the top of Mount Washington, 6,400 feet, to a distance of 800 miles, and from the highest mountain plateau on earth, 24,000 feet high, to a distance of 3,000 miles, almost one eighth of the circumference of the globe. Inversely, the tops of such mountains could be seen from similar distances, and every one knows that this is by no means the case. The fact is that the dip is only about 8 inches for the first mile; but for 2 miles it is nearly 2 feet, for 8 miles 4 feet, for 7 miles 30 feet, and for 11 miles not less than 88 feet. This is clearly shown in the engraving,



wherein the circle represents a section of the earth through its center, O. PI is the horizon of the point, P, and Rr, Ss, and Ii are heights from which the point, P, can be seen at different distances, Pr, Ps, and Pi. Without going into any mathematical demonstration, it is clear that the heights Rr, Ss, and Ii increase in a far greater ratio than the distances, Pr, Ps, and Pi; but in order to find the relation between the respective heights and distances correctly, a simple trigonometrical calculation is required, without which the solution of the problem is impossible.

ORP, OSP, and OIP are rectangular triangles, in which the angles at R, S, and I are the complements of those at O. Let the distance, PS, be 70 miles, about 1°; then the angle, PSO, will be 89°; and as the sides, PO and SO, of the triangle, PSO, are to one another in ratio as the radius to the sine of the angle, PSO, we will have: SO : PO = rad. : sin. 89°.

As PO is the radius of the earth, 20,891,914 feet, we will have: SO : 20,891,914 = 1 : 0.9998477, from which SO = 20,894,954. From this we subtract the earth's radius, Os, leaving 3,040 feet for Ss, the height required to see the point, P, at a distance of 70 miles.

In the same way, other distances may be calculated, and we have condensed some items of these calculations into the following table:

TABLE OF THE RELATION BETWEEN HEIGHTS AND DISTANCES, SEEN ON THE EARTH'S SURFACE.

For	1', nearly 1 mile, the height is	8 inches.
" 2', "	2 "	2 feet.
" 3', "	3 "	4 "
" 6', "	7 "	30 "
" 10', "	11 "	88 "
" 20', "	23 "	338 "
" 30', "	35 "	760 "
" 35', "	41 "	1,036 "
" 50', "	58 "	2,116 "
" 60', "	70 "	3,040 "
" 80', "	93 "	5,430 "

These results are proved by experiment to be correct, as we shall find when traveling in mountainous districts and noting how far we can see. At the highest tops of the Highlands, on the Hudson river, near West Point, which do not reach 2,000 feet, we can, on a clear day, just get a glimpse of the highest buildings in New York city, using a telescope. The distance is only fifty miles; but at a height of 1,600 feet, objects 50 miles off are invisible. In order to see to a distance of 90 miles, the height necessary is about 5,000 feet; and if the Philadelphia tower is built to a height of 760 feet, objects at a distance of 35 miles only, less than half way to New York, may be seen. If, however, two towers were built, one in New York and the other in Philadelphia, each 1,200 feet high, from each a circle of 45 miles radius would be visible, and the top of the one would be just perceptible from the top of the other, by means of a telescope, if the atmosphere were exceptionally clear. X.

To prevent water freezing in the gas meter add glycerin. The proper proportion is one pint of glycerin to a gallon of water.