

Scientific American.

ESTABLISHED 1845.

MUNN & CO., Editors and Proprietors.

PUBLISHED WEEKLY AT NO. 37 PARK ROW, NEW YORK.

A. D. MUNN. A. E. BEACH.

TERMS FOR THE SCIENTIFIC AMERICAN.

One copy, one year, postage included. \$3 20 One copy, six months, postage included. 1 60

Clubs.—One extra copy of THE SCIENTIFIC AMERICAN will be supplied gratis for every club of five subscribers at \$3.20 each; additional copies at same proportionate rate. Postage prepaid.

The Scientific American Supplement

is a distinct paper from the SCIENTIFIC AMERICAN. THE SUPPLEMENT is issued weekly; every number contains 16 octavo pages, with handsome cover, uniform in size with SCIENTIFIC AMERICAN. Terms of subscription for SUPPLEMENT, \$5.00 a year, postage paid, to subscribers. Single copies 10 cents. Sold by all news dealers throughout the country.

Combined Rates.—The SCIENTIFIC AMERICAN and SUPPLEMENT will be sent for one year, postage free, on receipt of seven dollars. Both papers to one address or different addresses, as desired.

The safest way to remit is by draft, postal order, or registered letter.

Address MUNN & CO., 37 Park Row, N. Y.

Subscriptions received and single copies of either paper sold by all the news agents.

Publishers' Notice to Mail Subscribers.

Mail subscribers will observe on the printed address of each paper the time for which they have prepaid. Before the time indicated expires, to insure a continuity of numbers, subscribers should remit for another year. For the convenience of the mail clerks, they will please also state when their subscriptions expire.

New subscriptions will be entered from the time the order is received; but the back numbers of either the SCIENTIFIC AMERICAN or the SCIENTIFIC AMERICAN SUPPLEMENT will be sent from January when desired. In this case, the subscription will date from the commencement of the volume, and the latter will be complete for preservation or binding.

VOL. XXXVII., No. 7. [NEW SERIES.] Thirty-second Year.

NEW YORK, SATURDAY, AUGUST 18, 1877.

Contents.

(Illustrated articles are marked with an asterisk.)

Table listing various articles such as American Institute Exhibition, Aniline colors and varnish, Antelope, the curly haired, Bad water, to detect, Base ball, curving, Blacking, boot (33), Carbolic soap (18), Carpet loom, the Gates, Cement for parchment paper (10), Cement for belts (7), Citric acid, to make (37), Cleopatra's needle, Coal fields of Ohio, Compass, good ship's, Concrete mixer, Correspondence, Washington, Davyrum, Dead bodies, preserving, Diphtheria, hypsulphuric in, Dustpan, improved, Electric fire, Electrical lamp, Ether, freezing point of, Expandibility of metals (49), Eyes, dimmed, Feed cutter, improved, Fertilizer from blood, Filtering sea water (1), Fish from artesian wells, Flower bed, floating, Fruit canned in Cincinnati, Glass gauges, to cut (31), Horse, speed of, pressure, etc. (56), Hose, india rubber, Ice machine, Holden's, Inventions patented in England, Iron, to galvanize (16), Jackson, Isaac W., Kid gloves, Kid gloves, to clean (20), Manganese (4), Marble, ink stains removed (5), Mars and Saturn, conjunction of, Measures, standard (56), Mercury, pressure of (55), New books and publications, Oxygen in the sun, Patents, American and foreign, Patents, official list of, Peroxide of hydrogen (19), Pigeon vs. locomotive, Poison ivy and its remedies, Railway strike, H. W. Beecher on, Rotation in motion, Shellac varnish (27), Shower, artificial, Silver first coined, Solar spectrum, new theory of, Starch, laundry (9), Stumps, to destroy (12), Sulphur in acetic acid, Sunshine in London and N. Y., Sulphurous odors, Tin from copper, to remove, Tobacco, to flavor (53), Tooth paste (32), Venus' slippers, Vinegar, sulphuric acid in, Wash boiler, improved, Watch, invention and history, Water snake, Wheel, revolution of (51), White metal (13), Woods, gravity of, Zinc dust, combustion of

TABLE OF CONTENTS OF

THE SCIENTIFIC AMERICAN SUPPLEMENT,

No. 85,

For the Week ending August 18, 1877.

I. ENGINEERING AND MECHANICS.—Eight Horse Traction Engine. With 4 illustrations.—The Disconnecting Compound Screw Engines of Steamship "Otter." With one page of engravings.—The New Fast Irish Mail Steamers.—Analysis of Boiler Incrustations. By EDWARD FRANCIS.—Improved Tug Steamers, with bow and stern screws.—The theoretical Steam Engine. By CHARLES E. EMERY.—The Boettius Gas Furnace.—Lath Drill Machine. Lath Drill Machine. With 1 engraving.—General Description of the New Water Works, Detroit, Mich.—Iron Railroad Ties as used on the Central Pacific Railway.—The Highest Railway Level in the United States.—The Chelsea Swimming Baths, London. With an engraving of the three baths.—The New Post Office and Custom House, Albany, N. Y. With an engraving.—On Ice-making Machines; by W. N. HARTLEY, F.R.S.E. Being a concise description of the best methods for the artificial production of ice.—On the Manufacture of Chinese Gongs; showing the method of production, the alloys used, etc.—How to Draw a Straight Line. By A. B. KEMPE, A.B. With 12 illustrations.—On Vortex Motion. By Professor OSBORNE REYNOLDS. An interesting and valuable paper.—Results of the Chronometer Trials at Geneva, 1876-77.—Discussion on the Parallelogram of Forces. By D. P. BLACKSTONE. II. ELECTRICITY, LIGHT, HEAT, ETC.—The Story of the Prism. A concise and interesting history of the Analysis of Light, and its marvelous results.—Action of Light upon Iodide of Potassium, Santele, and Eloeacca Oil.—New Method of Producing Light by the aid of Electricity.—New Experiment for showing the Composition of White Light.—On the Repair of Submarine Telegraph Cables.—Influence of Light on the Electrical Resistance of Metals.—Diathermancy of Rock Salt.—Suspension of Boiling Water on a Muslin Net.—Experimental Determination of the Thickness of Soap Films.—The World of Matter.—Temperature and Organic Remains in Tahoe and Echo Lakes. III. TECHNOLOGY.—On the Principles of Tapping.—On Arlonine Blacks.—On Garnets; selection of the woods and manipulation; Garnet Prepare; Dye Beck; Brazil Garnet; Garnet with Sanders and Orchil; second Bath; third Bath.—New harmless method of giving permanent Green Color to Peas and other Canned Vegetables.—New Lichen Acid.—Cremation of Street Refuse. IV. AGRICULTURE, HORTICULTURE.—On the Home Manufacture of Superphosphates.—Value and best methods of Curing and Using Clover.—On the best methods of Potato Culture.—Economic Plants in the Island of Jamaica. An interesting paper showing the success of efforts to introduce and acclimatize many useful and valuable plants; most of which may be grown in the United States. V. ASTRONOMY.—Dr. Dohereck's recent Catalogue of thirteen Double revolving Stars.—Physical Observations of the planet Mars.—Mars in the Autumn of 1877. By RICHARD A. PROCTOR, F.R.S. With Chart of her lands and waters of the planet and 6 illustrations of her telescopic appearance in opposition. VI. SCIENTIFIC AMERICAN CHESS RECORD.—Constitution of the American Chess and Problem Association.—Portrait of Samuel Loyd; notices of some of his successes as a Chess Player; with selections of his original problems from the recent Centennial Problem Tournament.—Game selected from the celebrated match played in London between Steinitz and Blackburn, 1876, prize £120; with notes thereon by Mr. Steinitz.

Remit by postal order. Address

MUNN & CO., PUBLISHERS,

37 Park Row, New York.

Single copies of any desired number of the SUPPLEMENT sent to any address on receipt of 10 cents.

POISON IVY AND ITS REMEDIES.

A correspondent asks: "Can you inform me by what characteristics I can determine the poisonous species of dogwood and ivy, and what simple remedy there is for their effect on the skin?"

Poisonous dogwood is a name improperly given in some parts of the United States to the rhus venenata, a species of poisonous sumach. It is sometimes called poison elder. It is a neat, graceful shrub growing from 6 to 18 feet high, and is found in swamps from Canada to Louisiana. The young shoots are purple or green clouded with purple, and marked by orange-colored dots which turn grayish; the leaves have 7 to 13 leaflets, which are dark green, pointed and entire on the margins; the greenish yellow flowers are in loose axillary panicles, and the greenish white fruit hangs in loose clusters on stems 6 to 8 inches long, and remains after the leaves have fallen; the juice is milky, and dries to a black varnish. This has poisonous qualities which are virulent. Its effect is an acute eczematous inflammation of the skin, often accompanied with much swelling.

The poison ivy or poison oak, in some places called mercury vine, the toxicodendron group of the botanists, includes two species with white or dun-colored berries in loose panicles and highly poisonous foliage. It has leaves of 3 leaflets, which are rhombic ovate, and variously notched, lobed, or even entire; its flowers are in loose slender axillary panicles; the smooth fruit is pale brown. It is found nearly all over the country, and especially in moist and shady places, and presents two forms, one erect and the other climbing. It clammers over rocks and fences, and by means of aerial rootlets ascends the trunks of the tallest trees, and adheres with great pertinacity. When wounded it exudes a milky juice, which becomes black on exposure to the air, and upon fabrics makes a stain indelible by all ordinary solvents. The leaves taken internally promote the secretions of the skin and kidneys. This plant is highly poisonous to some persons. Many can handle these plants without any unpleasant results, while others are seriously affected by touching them or even passing near them. The poisonous properties are due to a volatile acid, which has been called toxicodendric.

Many remedies are employed for poisoning by these plants, some of which will have beneficial effect on some persons, while on others have no effect at all. Water saturated with salt will often prove a cure, and at other times have no effect. The same may be said of sweet oil. There has been a remedy employed in some of the New England States that has been claimed to be effectual. It was this, the fat of the common black snake (bascanion constrictor) rendered into oil and applied to the parts affected. A strong lye made from wood ashes has been beneficially used, and so has an application of iodide of potassium. Another remedy is to take the fresh bark of the witchhazel (Virginian hamamelis), boil and apply the liquor as hot as the patients can bear it. A decoction made of the rattlesnake weed (hieracium venosum) applied to the parts afflicted will in most instances afford relief. Another remedy is to take one pint of the bark of the spotted alder, add one quart of water, and boil down to one pint; wash the parts poisoned several times a day. This remedy is said not to be injurious. Another remedy is to take the leaves of the poisonous nightshade (belladonna), boil them in milk to a poultice, bind it on the poisoned parts, and renew as often as it gets dry. A solution of belladonna, say a teaspoonful to a tumbler of water, with which bathe the parts freely. This has been used with signal success. Extract of lobelia or a poultice made from the fresh leaves may be used, but the external use of the plant in excess may produce vomiting and symptoms of poisoning. It ought to be applied under the advice of a physician. Another remedy is to bathe parts with spirits of niter. If the blisters are broken, so as to allow the niter to penetrate the cuticle, a simple application may effect a cure. Apply several times daily. Another remedy is to take three or four drops of the medicinal remedy known as rhus toxicodendron, drink two or three times daily in half a glass of water.

A NEW THEORY OF THE SOLAR SPECTRUM.

Since the invention of the spectroscopic, and its application to the study of the solar spectrum, the dark lines in the latter have been considered as absorption bands, caused by a layer of ignited metallic vapors, which surrounded the photosphere of the sun and changed the luminous and continuous spectrum of the photosphere into one covered with a multitude of dark lines, corresponding with the bright lines which we can produce by the combustion of various metallic substances.

Professor Henry Draper now comes forward with a series of experiments and deductions from the same, and proves that we must change this theory and form another conception, namely, that the solar spectrum consists also of bright lines and bands superposed on a less luminous background of continuous spectrum. Such a conception, combined with observations in regard to these bright lines, opens the way to the discovery of metalloids, sulphur, phosphorus, selenium, chlorine, bromine, iodine, fluorine, carbon, etc., the lines of which thus far have not been discovered in the solar spectrum. At the same time many of the dark lines, not thus far accounted for, may be due to being merely intervals between very bright lines.

That an incandescent gas in the solar atmosphere should not always be subject to the law, that it absorbs rays of the same refrangibility as it emits, may, at first sight, be difficult to understand. But the fact is, the substances thus far investigated in the sun have been metallic vapors, to which,

according to our present knowledge of chemistry, hydrogen also belongs. The metalloids may, and probably do, behave differently; the intensity of the light, from a great thickness of incandescent hydrogen, overpowers the effect of the photosphere; and instead of throwing a shadow of the rays of the same refrangibility, it increases the luminosity. It is as if a person looked through a yard thickness of ignited sodium vapor to a candle flame; he would see no dark sodium light, but a bright one; while looking at a very bright flame, he would see the comparatively dark sodium lines.

This would necessitate the supposition that some incandescent gases could give out more light than other substances in the sun, and why not? Has not Huggins shown that, in the outburst of the star r Coronae Borealis, hydrogen could give bright lines on a bright background of a similar nature to the background of the solar spectrum?

It is evident that bright lines on a less bright background make to ocular observations not so much impression upon the mind as the dark lines, and this is the simple reason that thus far they have been overlooked. If, however, the solar spectrum is photographed, such lines become very prominent; and the photograph being a permanent record, they may be easily compared with bright lines photographed from other spectra, such as those of air, oxygen, nitrogen, carbonic acid, etc., illuminated by means of the electric spark.

This is what Professor Henry Draper has been doing, and we call attention to the following article containing an account of the manner in which he demonstrated the presence of oxygen in the solar photosphere.

Fraunhofer who, about one century ago, first discovered the dark lines of the spectrum, which at the present day are named after him, also discovered that these lines are different when the light of some of the prominent fixed stars is investigated; and Berzelius, in remarking this, said in the beginning of this century that the study of these lines would at some future day lead us to the knowledge of the cause of the development of light in the heavenly bodies. This was a genuine prophecy, of which the world now begins to see the realization.

DISCOVERY OF OXYGEN IN THE SUN BY PHOTOGRAPHY.

Professor Henry Draper has announced the discovery of a series of bright lines or bands in the photograph of the solar spectrum, which correspond exactly with the principal bright lines or bands seen in photographs obtained by means of electric illumination in the spectrum of oxygen.

He has, in the American Journal of Science and Arts, published a paper and illustrated it with a photograph, in which he shows the perfect coincidence of certain bright lines. The photograph contains in its upper half the solar spectrum, and in its lower half the spectrum of air obtained by passing the spark of a Gramme induction machine (driven by Brayton's petroleum motor) from an iron to an aluminum point. The coincidence of the luminous oxygen and even of the nitrogen lines is really remarkable; and as the photograph is stated to be absolutely free from hand work or retouching, it places the subject in question beyond doubt. Thus the iron and aluminum lines, produced by the effect of the powerful electric current upon the electrodes, show themselves, and the first may be traced in the solar spectrum at the corresponding places, as might be expected.

We will only add that Professor Draper has made detailed comparison of these lines in the spectra of air, oxygen, nitrogen, hydrogen, carbonic acid, carburetted hydrogen, and cyanogen, so as to be sure of the luminous lines belonging to oxygen, and he has also made experiments with these gases at various pressures, as in some of them the lines vary with the pressure. It may be remarked as an important fact that the spectrum of oxygen is not subject to variation, but that its lines are constant at all pressures.

Science is already largely indebted to Professor Draper for the originality of his researches, and no doubt important results may be expected in the train of research he is now following. It is useless to speculate as yet on the nature of the sun, and it is better left to later times, when our knowledge of this remarkable body will be more complete; but one thing is certain, that the idea of Herschel that the sun may be an inhabited globe must be given up. It is undoubtedly a body at a temperature so high that the substances present there are dissociated and cannot enter into chemical combinations. However, that we will find there all the elements present on our globe may be anticipated if we adopt the theory of Kant and Laplace of a common origin of our whole planetary system out of one single nebula.

AN ELECTRIC FIRE.

A fire recently occurred at the Western Union Telegraph Office, in New York city, that was one of those incidental circumstances in the operation of a great enterprise that imparts a lesson of experience. The cause was defective insulation of wires that came in contact, in what is known as the "grand switch." This switch is situated in an upper story, and consists of a mahogany table about 25 feet long and 5 broad. It is of elegant cabinet work, placed vertically, and contains about 400 wires, which pass from the battery room through apertures in the ceiling into the switch. It also controls about 10,000 connections. It is, in fact, a systematized combination of all the wires issuing from the chief office to every part of the country.

These wires as they enter the switch are separated and insulated. By some means two of the wires, not sufficiently insulated, came in contact with each other. Electrical heat