explosive is to be exploded electrically by reversing the current. The steering is also done electrically through the agency of a polarized relay.

Its operation from a shore station is limited in extent by the length of the connecting cable. In the large torpedoes this will be two miles. In operation from a war ship, it is proposed to run two or more boats along a parallel course with the ship and close to it, receiving their motive power from the ship's electric plant. Here, as the ship and torpedoes will all progress alike, and as the electric plant on board can supply power for an indefinite period, the ship and torpedoes can keep company for almost any number of miles. When the enemy is approached the course of the ship can be arrested or its speed can be slackened, and the torpedo sent ahead or to either side at high speed, in order to destroy the enemy. For fort use a special form of casemate with conning tower and other necessary features have been planned by the company. There is little question that a number of these torpedoes could do much to defend New York harbor, at the end of the East River, between Willets Point and Fort Schuyler, and at the Narrows, from entrance by hostile vessels. Our thanks are due to Mr. Everett Frazar, the president of the company, for courtesies received.

# <u>. . . .</u>

A Liquid Volatile Nickel Compound. Nickel seems destined to startle the modern chemical world. From being a comparative rarity, except on plated goods, it became a common laboratory material but a little while ago, then its title to be considered an element was impugned, and now it is both triumphantly reinstated and its vapor density, for the first time, determined. These results have followed from the researches of Mr. Mond and Drs. Quincke and Langer, which were recently made the subject of a paper before the Chemical Society. The investigation owed its origin to the need for removing carbonic oxide

from producer gas, in order to use it in the gas battery described in Mr. Mond's presidential address at the annual meeting of the Society of Chemical Industry last year, nickel and cobalt being found to effect this object.

Working in this direction, it has been found that a direct compound of nickel and carbonic oxide, viz., Ni(CO)<sub>4</sub>, exists, the new substance being a colorless *liquid*, volatile at ordinary temperatures, boiling at 43° C., having a specific gravity of 1.3185 at 17° C., soluble in alcohol, benzine, and chloroform, and not acted upon by dilute acids and alkalies. Its vapor explodes when heated to 60° C. Its vapor density determined by Victor Meyer's method is 6'01, instead of the theoretical value, 5.9; from this, the atomic weight of nickel is found to be 58. The metal itself can be obtained from it in the form of brilliant metallic mirrors of such purity as to form a splendid raw material for value obtained by reducing nickel oxide from this source by heating in a stream of electrolytic hydrogen was 58.61, corresponding closely with that previously accepted, viz., 58.52. This proves conclusively that Kruss and Schmidt's assertion that the metal hitherto considered to be pure nickel is contaminated with another element, and that all data concerning it consequently need revision, cannot be sustained, and replaces nickel as we have always known it among the elements.

The constitution of the compound Ni(CO)<sub>4</sub> is still the subject of the keenest speculation. Mr. Mond declined to be "drawn" on this point, admitting the temptation to represent it by some fascinating ring formula, but contenting himself at present with a statement of the facts. In connection with this curious body, it is to be noted that no similar cobalt compound can be obtained, thus establishing another method of differentiation and separation between it and its twin brother nickel.

#### ..... Interesting and Useful Books.

## Several years ago the writer, traveling abroad, visited a monastery, and among other curiosities shown

Scientific American. ESTABLISHED 1845.

MUNN & CO., Editors and Proprietors. PUBLISHED WEEKLY AT

No. 361 BROADWAY, NEW YORK.

A. E. BEACH.

O. D. MUNN.

TERMS FOR THE SCIENTIFIC AMERICAN. 

# The Scientific American Supplement

the Scientific American Supplement is a distinct paper from the SciEntific American. Supplement is issued weekly. Kvery number contains 16 octavo pages, uniform in size with SciEntific American. Terms of subscription for SUPPLEMENT, \$5.00 a year, for U. S., Canada or Mexico. \$6.00 a year to foreign countries belonging to the Postal Union. Single copies, 10 cents. Sold by all newsdealers throughout the country. See prospectus last page. Comblord Kates.—The SCIENTIFIC AMERICAN and SOPPLEMENT will be sent for one year, to any address in U. S., Canada or Mexico, on receipt of seven dollars. To foreign countries within Postal Union, nine dollars a year. lollars a year.

### Building Edition.

Building Edition. THE ARCHITECTS AND BUILDERS EDITION OF THE SCIENTIFIC AMER-ICAN is a large and splendid illustrated periodical, issued monthly, con-taining floor plans, perspective views, and sheets of constructive details, pertaining to modern architecture. Each number is illustrated with beautiful plates, showing desirable dwellings, public buildings and archi-tectural work in great variety. To builders and all who contemplate build-ing this work is invaluable. Has the largest circulation of any architec-tural publication in the world. Single copies 25 cents. By mail, to any part of the United States, Canada or Mexico, \$2.50 a year. To foreign Postal Union countries, \$5.00 a year; Combined rate for BUILDING EDITION with SCIENTIFIC AMERICAN, \$5.00 a year; combined rate for BUILDING EDITION, SCIENTIFIC AMERICAN and SUPPLEMENT, \$3.00 a year. To foreign countries, \$1.50 a year.

#### Spanish Edition of the Scientific Americau.

Spanish Edition of the Scientific Americau. LA AMERICA CIENTIFICA E INDUSTRIAL (Spanish trade edition of the SCIENTIFIC AMERICAN) is published monthly, uniform in size and typo-graphy with the SCIENTIFIC AMERICAN. Every number of La America is profusely illustrated. It is the finest scientific, industrial trade paper printed in the Spanish language. It circulates throughout Cuba, the West Indies, Mexico, Central and South America, Spain and Spanish posses-sions-wherever the Spanish language is spoken. 3.00 space, post paid to any part of the world. Single copies 25 cents. See prospectus. MUNN & CO., Publishers, 361 Broadway, New York. 363 Broadway, New York.

361 Broadway, New York. praft or bank check. Make all remittances payable to order of MUNN & CO.



51

## TABLE OF CONTENTS OF SCIENTIFIC AMERICAN SUPPLEMENT No. 760.

#### For the Week Ending July 26, 1890.

Price 10 cents. For sale by all newsdealers PAGE

- I. BIOGRAPHY.-Miss Fawcett.-Portrait of and note on Miss Faw-cett, graded above the senior wrangler in Cambridge University.. 12142 II. BIOLOGY.-Protective Mimicry in a Fish.-The curious little sea horse.-Its resemblance to the plants among which it lives.-1il-
- Paste Eels.—Note on identity of vinegar eels and paste eels..... 12146 Paste Eels.-Note on identity of vinegar eels and paste eels..... 111. CHEMISTRY.-The Quantitative & Rstimation of Coloring Matters by Means of their Absorption Spectra.-By T. L. PATTERSON.-An interesting paper, indicating a method of spectroscopic de-termination of quantities.-With full description and illustrations of apparatus employed.-4 illustrations UV. CIVII. ENGINE EERING.-The Railways of India.-A picturesque account of the achievements of the civil engineer in the empire of India, with map of railroad lines.-1 illustration U. ELECTRICITY.-A Very Simple Rheostat.-A home-made piece of electrical apparatus for changing the intensity of a current.-1 illustration
- 12148
- 12136

#### IT PAYS TO ADVERTISE IN THE SCIENTIFIC AMERICAN.

In a letter to the publishers of this paper G. M. Robinson, Esq., president of the Charter Gas Engine Company, Sterling, Ill., writes that they are overwhelmed with orders, and that "inquiries from the SCIENTIFIC AMERICAN advertisement are so numerous and from so many points, both foreign and domestic, that we often wonder if there will ever be an end to them."

#### THE GYPSY MOTH IN MASSACHUSETTS.

The ravages of the gypsy moth (Ocneria dispar) in Massachusetts have been of such a very serious nature that, in March last, the legislature passed a bill creating a commission of three members, who were called upon by the terms of the act to carry into execution all possible and reasonable means and measures to prevent the spreading and to secure the extermination of the pest. Twenty-five thousand dollars was appropriated to defray the expense of the commission, and this sum has been increased by a subsequent appropriation to \$50,000.

In June, 1889, several caterpillars were sent to the Hatch experiment station, at Amherst, which is a part of the Massachusetts Agricultural College. Prof. C. H. Fernald, of the Division of Entomology, was absent in Europe, but Mrs. Fernald, who had charge of the entomological work during her husband's absence, immediately designated the insect as the gypsy moth (Ocneria dispar, Linn.) of Europe. In November, 1889, Prof. Fernald issued a special bulletin in which the following interesting facts regarding the history of the moth appear.

The gypsy moth is abundant in nearly all parts of Europe, Northern and Western Asia, and it even extends as far as Japan. In 1817 the cork oaks of Southern France suffered severely from the attacks of this insect. One of the papers of that time stated that the beautiful cork oaks which extended from Barboste to the city of Podenas were nearly destroyed by the caterpillars of the gypsy moth. After having devoured the leaves and young acorns, they attacked the fields of corn and millet and also the grape lands and fruit trees. In 1878 the plane trees of the public promenades of Lyons were nearly ruined by the same insect. In regard to his personal observations in Europe, Prof. Fernald says: "Only last summer I saw the moths in immense numbers on the trees in the Zoological Gardens in Berlin, where the caterpillars had done great injury; and the European works on entomology abound with instances of the destructiveness of this insect."

Authorities seem to agree in stating that the introduction of the gypsy moth into this country was accidental. In 1870, Mr. L. Trouvelat, a Frenchman residing at Medford, was experimenting with silkworms, and a few of these caterpillars were among the silkworms, and some of them escaped. It is a curious fact that, although the insect has been in this country since 1870, it did not become a pest until last season, and according to the entomologists of the United States Department of Agriculture, had not, up to about March 1, found its way into the collections or been mentioned in the check lists.

One of the first acts of the commissioners appointed to conduct the campaign against the gypsy moth was to advertise for specimens, for the purpose of ascertaining if they were to be found in any other portion of the State than about Medford and Malden. A number of specimens were received, but none proved to be the gypsy moth. It is therefore deemed to be certain that they are only found in this country in the same region where they were originally liberated twenty years ago. The section which has been devastated by them extended last season three miles in length and one in width, but this season the territory has been extended until it is fifteen miles long and four wide.

As the depredations of this insect and the means which are being used to check them are attracting wide attention, the following minute description by Prof. Fernald in its various stages of development

him was a series of bound volumes, the sides of which were made of polished boards from the forests of the country, showing the grain of the woods.

At first sight the volumes presented the aspect of bundles of wood. But after a more careful examination it was found that they contain a detailed account of the trees that they represent. On the back, the bark has been detached in order to describe the title of the book by its scientific and its common names.

One of the pages is formed by a broken piece of the wood of the tree, showing its fibers and natural fractures; the other shows the wood when it has been polished and varnished. At one of the ends the fibers are seen as they remain after the passage of the saw, and at the other, the wood finely polished. Upon opening the book, the fruit, the grain, the leafage and other productions of the tree, the moss which generally grows on the trunk, and the insects which live on its different parts are seen. Added to this is a well printed description of the habits of the tree, the places where it grows, and its method of growth.

12143 12144 12135 maceutical art, a good prognosis for future progress. XIII. T KCHNOLOGY.—Cane Cultureand Sugar Manufacturein Cuba. —A Southern industry, the cost and economies of cultivature the sugar cane and manufacturing the sugar. Lighting by Compressed Gas.—The use of compressed gas for maintenance of illuminated buoys, the plant for its production and buoys from which it is burned.—S illustrations.... Pick's System of Manufacturing Salt in Vacuo.—Theapplication of the triple effect apparatus and the well known strike pan of the sugar manufacturer to the salt process, full details of apparatus and process.—1 illustration. 12145 1214 12140 12138

may be of interest, especially as the vigorous measures which have been adopted in Massachusetts may result in driving it to other localities.

This insect was first described by Linneus in 1758, in the tenth edition of his Systema Naturæ, Vol. 1, page 501, under the name of  $Bombyx \ dispar$ , and while it has retained the specific name of dispar, the European entomologists, since the time of Linneus, have given several different generic names, as Liparis, Hypogymna, Porthetria, Ocneria and Psilura, but I have adopted that given by Staudinger in his Catalogue of the Lepidoptera of Europe-Ocneria dispar.

Several different common names have also been given to it in Europe, as the sponge moth, the gypsy moth, the great-headed moth, the fungus moth and others, but I have adopted the one used by the English entomologists-the gypsy moth.

The males are of a yellowish brown color, with two dark brown lines crossing the forewings, one at the basal third, the other on the outer third, somewhat curved, and with teeth pointing outward on the veins,